

## Hypertension

Rationale for drug use | BP measurement | Before starting drug treatment | BP targets | Drug treatment

### Cardiovascular drugs

- Drugs for heart failure
- Drugs for angina
- Antihypertensives
  - Hypertension
  - Thiazide and related diuretics
  - Other diuretics
  - ACE inhibitors
  - Sartans
  - Calcium channel blockers
  - Beta-blockers
  - Other antihypertensives
- Drugs for arrhythmias
- Drugs for dyslipidaemia
- Drugs for pulmonary hypertension
- Drugs for other cardiovascular disorders
- Tables
  - Management of heart failure with reduced ejection fraction
  - Comparative information for sympathomimetics
  - Comparative information for nitrates
  - Comorbidities affecting antihypertensive choice
  - Antihypertensive fixed-dose combinations
  - Comparison of calcium channel blockers
  - Comparison of oral beta-blockers
  - Drugs that may prolong QT interval
  - CHA<sub>2</sub>DS<sub>2</sub>-VA score
  - Comparison of lipid-lowering drugs

### Hypertension

See also [www.heartfoundation.org.au/conditions/hypertension](http://www.heartfoundation.org.au/conditions/hypertension)<sup>12</sup>, [www.heartfoundation.org.au/Conditions/EP-Absolute-CVD-Risk-Clinical-Guidelines](http://www.heartfoundation.org.au/Conditions/EP-Absolute-CVD-Risk-Clinical-Guidelines)<sup>12</sup>

Observational studies show that the risk of cardiovascular morbidity and mortality rises as BP increases from 115/75 mm Hg; age, sex (male) and other cardiovascular risk factors contribute to this risk.

The distinction between normotension and hypertension is arbitrary: generally normotension is considered as clinic BP of 120 mm Hg systolic (SBP)/80 mm Hg diastolic (DBP) and below, and hypertension as >140/90 mm Hg.

The decision to start treatment should incorporate thorough assessment of BP, overall cardiovascular risk and end-organ damage.

#### Rationale for drug use

Reduce premature cardiovascular morbidity and mortality and microvascular disease affecting the brain, kidneys and retinas.

#### BP measurement

Confirm level of BP with multiple readings on different occasions; ensure correct cuff size; consider possibility of white coat effect on BP.

Measurement method affects the results: ambulatory and home BP are generally lower than clinic BP and are stronger predictor of outcomes.

Automated office BP measurement, which may be taken unobserved by staff, provides substantially lower SBP (by about 10–20 mm Hg) than conventional clinic readings. This method was used in the SPRINT study but is currently not often practised in Australia.

#### Before starting drug treatment

Look for secondary causes of hypertension (including drugs, eg NSAIDs, corticosteroids) and evidence of end-organ damage.

Identify and treat other modifiable cardiovascular risk factors, such as obesity and dyslipidaemia; encourage smoking cessation.

Encourage a healthy lifestyle for all patients (eg weight loss, increased physical activity, healthy diet). Such interventions can reduce BP (potentially avoiding or delaying the need for antihypertensives). Regular aerobic exercise can reduce daytime SBP by up to 3.2 mm Hg, and a weight loss of 10 kg in an overweight person may reduce SBP by 6–10 mm Hg.

#### Assess cardiovascular risk

Assess absolute cardiovascular risk (probability of an event within 5 years) for adults without cardiovascular disease aged 45 years and older (35 years for Aboriginal and Torres Strait Islander people), using the National Vascular Disease Prevention Alliance tools:

- risk charts at [www.heartfoundation.org.au/health-professional-tools/cvd-risk-charts](http://www.heartfoundation.org.au/health-professional-tools/cvd-risk-charts)<sup>12</sup>
- risk calculator at [www.cvdcheck.org.au](http://www.cvdcheck.org.au)<sup>12</sup>.

Cardiovascular risk calculators use clinic BP readings. Use of home, ambulatory or automated office BP in a risk calculator may inappropriately underestimate risk.

Some patients may have additional risk factors (not considered in the calculator), eg family history.

A risk estimate is not needed for patients already considered high risk, eg:

- diabetes and >60 years of age or with microalbuminuria
- moderate or severe chronic kidney disease (eg eGFR <45 mL/minute/1.73 m<sup>2</sup> or macroalbuminuria)
- established vascular disease
- familial hypercholesterolaemia; serum total cholesterol >7.5 mmol/L
- SBP >180 mm Hg or DBP >110 mm Hg
- Aboriginal and Torres Strait Islander people >74 years.

#### Management with consideration of cardiovascular risk

Provide lifestyle advice for all patients regardless of cardiovascular risk or drug therapy, offering ongoing support and follow-up to maintain long-term lifestyle changes.

##### Low risk (<10%)

Start antihypertensive treatment if persistently elevated SBP >160 mm Hg or DBP >100 mm Hg.

##### Moderate (10–15%) to high risk (>15%)

Start antihypertensive treatment if persistently elevated SBP >140 mm Hg or DBP >90 mm Hg.

#### BP targets

Australian guidelines recommend treating patients with uncomplicated hypertension to a target of <140/90 mm Hg or lower if tolerated.

In practice, BP can be lowered as far as it is tolerable; even if targets are not reached, any reduction in BP reduces risk of cardiovascular morbidity and mortality.

A lower target of SBP <120 mm Hg can be considered in selected high cardiovascular risk populations, provided there is close monitoring for adverse effects (eg hypotension, electrolyte abnormalities).

In the SPRINT study, patients at high cardiovascular risk showed fewer cardiovascular events and reduced mortality (but more treatment-related adverse events) when the target SBP was <120 mm Hg compared with <140 mm Hg. However, its generalisability and application to practice may be limited by its use of automated office BP measurement (see [BP measurement](#)) and exclusion criteria (eg age <50 years, diabetes, heart failure, history of stroke).

#### Drug treatment

Generally start treatment with a single drug at the lowest recommended dose (this will satisfactorily reduce BP in about 25–50% of people).

The presence of coexisting conditions may affect the choice of antihypertensive ([Table](#)).

For uncomplicated hypertension, unless there is a contraindication or a specific indication for another drug, first consider:

- an ACE inhibitor (or sartan) or
- a dihydropyridine calcium channel blocker or
- if 65 or older, a thiazide diuretic (low dosage).

Choose an agent given once daily.

Generally allow 4–6 weeks to assess response to treatment. For those with marked hypertension or high cardiovascular risk, consider a shorter review period.

Not all patients respond to all drugs; monitor response carefully; if initial drug not tolerated or has no effect, change to a drug from a different class.

**Table – Comorbidities affecting antihypertensive choice**

Comorbidity	Drugs with favourable effect
chronic kidney disease with micro- or macroalbuminuria	ACE inhibitors (or sartans)
heart failure	ACE inhibitors (or sartans), beta-blockers (carvedilol, controlled release metoprolol, bisoprolol, nebivolol), thiazide diuretics
post MI	beta-blockers (except pindolol), ACE inhibitors (or sartans)
angina	beta-blockers (except pindolol), calcium channel blockers, ACE inhibitors
AF	ACE inhibitors or sartans (verapamil, diltiazem, beta-blockers may help rate control)
Comorbidity	Drugs with unfavourable effect
asthma, COPD	beta-blockers <sup>1</sup>
bradycardia, second- or third-degree atrioventricular block	beta-blockers, diltiazem, verapamil
heart failure	calcium channel blockers (especially verapamil, diltiazem)
gout	thiazide diuretics
<sup>1</sup> cardioselective beta-blockers (eg metoprolol) may be used cautiously in COPD or well-controlled asthma, see <a href="#">Respiratory</a> in Beta-blockers	

#### Inadequate effect

Add a second antihypertensive rather than increasing the dose of the first (which may cause adverse effects without improving BP control).

The preferred combinations are:

- ACE inhibitor (or sartan) with a calcium channel blocker or a thiazide diuretic
- calcium channel blocker with a thiazide diuretic.

Other effective combinations include a beta-blocker with an ACE inhibitor (or sartan) or a dihydropyridine calcium channel blocker. Use of beta-blockers with thiazide diuretics is also effective (but increases risk of diabetes compared with other combinations). Fixed-dose combinations may be convenient and less expensive, see [Table – Antihypertensive fixed-dose combinations](#).

Unless there are specific benefits, avoid:

- diltiazem or verapamil with beta-blocker (risk of severe bradycardia and heart block)
- ACE inhibitor (or sartan) with potassium-sparing diuretic (risk of hyperkalaemia)
- ACE inhibitor with sartan (benefit over ACE inhibitor alone is questionable, may increase renal complications; specialist use only, see [Practice points](#) in Sartans).

#### BP target still not reached

If both drugs are well tolerated, increase dose of one agent (non-thiazide) towards the maximum recommended; if this is still inadequate, increase the dose of the other antihypertensive.

If BP remains above the target despite maximum doses of at least 2 appropriate agents, consider other factors, eg poor compliance, high salt intake, secondary hypertension (including drug-induced), volume overload, sleep apnoea (can cause treatment resistance).

Three or more antihypertensives from different classes may be needed.

Resistant hypertension, which is uncontrolled BP despite 3 or more optimally tolerated antihypertensives (including a diuretic), occurs in >10% of patients. Consider primary aldosteronism especially if hypokalaemia is present. Seek specialist advice after ruling out compliance issues and other secondary causes.

[Spironolactone](#) is an effective add-on drug for resistant hypertension uncontrolled by a combination of first-line agents (note: risk of hyperkalaemia is increased when combined with ACE inhibitors or sartans particularly in the presence of kidney disease).

**Table – Antihypertensive fixed-dose combinations**

Brand <sup>§1</sup>	ACE inhibitor	Calcium channel blocker <sup>2</sup>	Sartan	Thiazide or related diuretic	Potassium-sparing diuretic	Statin
Abisart HCT			irbesartan	hydrochlorothiazide		
Accuretic	quinapril			hydrochlorothiazide		
Adesan HCT			candesartan	hydrochlorothiazide		
Asartan HCT			candesartan	hydrochlorothiazide		
Atacand Plus			candesartan	hydrochlorothiazide		
Avapro HCT			irbesartan	hydrochlorothiazide		
Avasartan HCT			irbesartan	hydrochlorothiazide		
Cadvast		amlodipine				atorvastatin
Caduet		amlodipine				atorvastatin
Candesan Combi			candesartan	hydrochlorothiazide		
Co-Diovan			valsartan	hydrochlorothiazide		
Coveram	perindopril arginine	amlodipine				
Coversyl Plus	perindopril arginine			indapamide		
Dilatart HCT			valsartan	hydrochlorothiazide		
Exforge		amlodipine	valsartan			
Exforge HCT		amlodipine	valsartan	hydrochlorothiazide		
Fosetic	fosinopril			hydrochlorothiazide		
Idaprex Combi	perindopril erbumine			indapamide		
Indosyl Combi	perindopril erbumine			indapamide		
Karvezide			irbesartan	hydrochlorothiazide		
Micardis Plus			telmisartan	hydrochlorothiazide		
Mizart HCT			telmisartan	hydrochlorothiazide		
Moduretic				hydrochlorothiazide		amiloride
Olmekar		amlodipine	olmesartan			
Olmekar HCT		amlodipine	olmesartan	hydrochlorothiazide		
Olmertan Combi			olmesartan	hydrochlorothiazide		
Olimetec Plus			olmesartan	hydrochlorothiazide		
Perindo Combi	perindopril erbumine			indapamide		
Prexum Combi	perindopril arginine			indapamide		
Pritor/Amlodipine		amlodipine	telmisartan			
Raptan	perindopril arginine	amlodipine				
Renitec Plus	enalapril			hydrochlorothiazide		
Sevikar		amlodipine	olmesartan			
Sevikar HCT		amlodipine	olmesartan	hydrochlorothiazide		
Tarka	trandolapril	verapamil				
Teltartan HCT			telmisartan	hydrochlorothiazide		
Tevetan Plus			srosartan	hydrochlorothiazide		
Triasyn	ramipril	felodipine				
Twynsta		amlodipine	telmisartan			
Zan-Extra	enalapril	lercanidipine				

<sup>1</sup> generic brands that name the drugs in the brand name may also be available

<sup>2</sup> all, except verapamil, are dihydropyridines

#### Drug choice

See also [Table – Comorbidities affecting antihypertensive choice](#)

It is the reduction in BP that is most important for reducing cardiovascular events, rather than the antihypertensives chosen. ACE inhibitors, calcium channel blockers, sartans and thiazide diuretics are usually used.

#### ACE inhibitors

First-line treatment, especially in patients with chronic kidney disease, diabetes with micro- or macroalbuminuria, is heart failure or with left ventricular dysfunction, in particular following MI.

#### Sartans

May be used as an alternative first-line treatment especially in patients who are intolerant of ACE inhibitors.

#### Calcium channel blockers

The dihydropyridines (amlodipine, felodipine, lercanidipine, nifedipine) are suitable as first-line treatment.

In meta-analyses, they reduced the frequency of stroke but increased rates of heart failure compared with other antihypertensive classes, while the incidence of cardiovascular events and mortality were similar.

Verapamil and diltiazem are contraindicated for those who also have heart failure, while dihydropyridines may be used cautiously.

#### Thiazide diuretics

Are generally well tolerated and are a first-line treatment for hypertension in those aged >65 years. Due to their association with new-onset diabetes, they are no longer recommended as first-line monotherapy in younger patients.

#### Beta-blockers

Compared with other antihypertensive classes, they have less effect in reducing the incidence of stroke and are associated with an increased risk of diabetes.

They are recommended for hypertension when compelling indications, such as angina, are present, see [Table – Comorbidities affecting antihypertensive choice](#).

#### Other drugs

A number of other drugs, such as the selective alpha-blockers, are still used but there is no reliable evidence from randomised trials about their effects on cardiovascular morbidity or mortality.

#### Special cases

##### Elderly

Start treatment with the lowest dose and titrate slowly.

In a subgroup of SPRINT participants, ie selected ambulatory patients aged >75 with hypertension (excluding comorbidities, eg diabetes, heart failure, history of stroke), lowering SBP to <120 mm Hg significantly reduced the risk of cardiovascular and all-cause mortality. Note that this study used automated office BP (where a lower SBP target of <120 mm Hg is comparable to clinic SBP <130–140 mm Hg), see [BP measurement](#).

Aim for the lower BP targets only when treatment is well tolerated and with close monitoring (eg for orthostatic hypotension, electrolyte abnormalities, acute kidney injury).

##### Coronary artery disease

Monitor BP closely when using antihypertensives. Both high (SBP >140 mm Hg or DBP >80 mm Hg) and low (SBP <120 mm Hg or DBP <70 mm Hg) BP were associated with increased risk of cardiovascular morbidity and mortality in hypertensive patients with increased risk of cardiovascular disease in an international cohort study.

##### Diabetes

BP target <140/90 mm Hg is recommended. If preventing stroke is a priority, consider lowering to a SBP <120 mm Hg, provided the patient can be closely monitored for adverse effects, eg hypotension, acute kidney injury.

Controlling BP improves cardiovascular morbidity and mortality and reduces the progression of diabetic nephropathy in people with both hypertension and diabetes.

ACE inhibitors or sartans are preferred in diabetes as they delay progression of renal disease in patients with micro- or macroalbuminuria.

Calcium channel blockers, thiazide diuretics and beta-blockers may also be used, however, although thiazide diuretics and beta-blockers have beneficial effects on cardiovascular outcomes in diabetics, they tend to worsen glycaemic control compared with other classes.

##### Hypertensive crisis

Avoid precipitous BP reduction (>25% over 2 hours) to prevent serious events, eg MI, stroke. Monitor BP continuously.

Seek specialist advice. Aim is to limit acute end organ damage.

Severe pre-eclampsia, rarely cerebral haemorrhage). Where there is no end organ damage, gradual BP reduction over several days is sufficient and safer.

To reduce BP over 2–6 hours, consider agents such as IV labetalol, sodium nitroprusside, glyceryl trinitrate, hydralazine and clevidipine. Beta-blockers may be needed to control reflex tachycardia. In most cases, aim towards 160/100 mm Hg rather than normal BP levels; then wait 24 hours before lowering further.

For less urgent BP reduction, use oral antihypertensive drugs and follow up within 24–72 hours.

##### Hypertension in pregnancy

For further information see [somanz.org/guidelines.asp](http://somanz.org/guidelines.asp)<sup>12</sup>.

BP targets and thresholds at which to begin drug treatment differ from those for non-pregnant women.

*Mild-to-moderate hypertension:* first-line drugs are methyl dopa and labetalol; nifedipine, hydralazine or prazosin may be used second line. Diuretics and other beta-blockers are generally avoided during pregnancy. ACE inhibitors and sartans must not be used in the second and third trimesters.

Severe hypertension: see local protocols and also [Pre-eclampsia and eclampsia](#). Oral nifedipine (conventional tablet, available through the SAS) and parenteral drugs, eg labetalol and hydralazine, are used for urgent BP reduction.